

USING ^{18}O TO PARTITION ECOSYSTEM CARBON EXCHANGES: IMPACT OF THE NEAR-SURFACE $\delta^{18}\text{O}$ VALUE OF SOIL WATER ON THE $\delta^{18}\text{O}$ VALUE OF THE SOIL-SURFACE CO_2 FLUX

William J. Riley

Contact: 510/486-5036, [wj Riley@lbl.gov](mailto:wjriley@lbl.gov)

RESEARCH OBJECTIVES

The ^{18}O content of atmospheric CO_2 has been proposed as a means to partition site-level measured net ecosystem carbon fluxes into component gross fluxes and, at the global scale, to estimate the regional distribution of CO_2 fluxes. However, these approaches require accurate prediction of the $\delta^{18}\text{O}$ value of the soil-surface CO_2 flux (δF_s). This work aims to better characterize and to improve the computational efficiency of models used in global and regional simulations.




Figure 1 (a) δF_s as predicted by ISOLM and the HDMR approach. Also shown is the gradient in δ_{sw} over the top 15 cm. (b) Cumulative isoflux from the soil to the atmosphere for the two approaches. The error the three-month growing season was 0.2%.

APPROACH

The net C^{18}OO surface flux depends, nonlinearly, on the depth-dependent $\delta^{18}\text{O}$ value of soil water (δ_{sw}), soil moisture and temperature, soil CO_2 production, and the $\delta^{18}\text{O}$ value of above-surface CO_2 (Riley, 2003a). We developed ISOLSM (Riley et al., 2002; Riley et al., 2003b) to simulate these processes within an established land surface model (LSM1). ISOLSM simulates the ^{18}O content of canopy water vapor, leaf water, and vertically resolved soil water; leaf photosynthetic C^{18}OO fluxes; CO_2 oxygen isotope exchanges with soil and leaf water; soil CO_2 and C^{18}OO diffusive fluxes (including abiotic soil exchange); and ecosystem exchange of H_2^{18}O and C^{18}OO with the atmosphere. Since ISOLSM is a computationally expensive model, we applied a high-dimension model representation (HDMR) technique to efficiently predict δF_s .

ACCOMPLISHMENTS

Our results indicate that the HDMR approach is very accurate and about 100 times faster than the full numerical solution of the C^{18}OO surface flux, making it appropriate for regional and global simulations. We successfully tested the HDMR approach over a growing season at a C_4 -dominated tallgrass prairie site, and then used the model to investigate the factors important in determining δF_s . The top panel of the figure shows comparisons between the full numerical model and the HDMR approach for a 20-day period. The bottom panel shows comparisons of the cumulative isoflux from the soil surface over the full season; the error over the growing season was less than 0.2%. The largest changes in δF_s occur when gradients in the top 5 cm are large. These conditions typically occur when soil evaporation is large, i.e., following precipitation.

SIGNIFICANCE OF FINDINGS

Simulation results indicate that δF_s is dependent on the $\delta^{18}\text{O}$ value of soil water in the top few centimeters of soil. These results indicate that recent approaches to estimating global distributions of the surface C^{18}OO flux are problematic and demonstrate the importance of accurately resolving near-surface δ_{sw} . Also, the development of the HDMR approach allows for accurate and computationally affordable simulations of regional and global distributions.

RELATED PUBLICATIONS

- Riley, W.J., Impact of the near-surface $\delta^{18}\text{O}$ value of soil water on the $\delta^{18}\text{O}$ value of the soil-surface CO_2 flux: Application of a high-dimension model representation technique. GRL, 2003a (submitted).
- Riley, W.J., C.J. Still, M.S. Torn, and J.A. Berry, A mechanistic model of H_2^{18}O and C^{18}OO fluxes between ecosystems and the atmosphere: Model description and sensitivity analyses. Global Biogeochemical Cycles, 16, 1095–1109, 2002.
- Riley, W.J., C.S. Still, B.R. Helliker, M. Ribas-Carbo, S. Verma, and J.A. Berry, Measured and modeled $\delta^{18}\text{O}$ in CO_2 and H_2O above a tallgrass prairie. Global Change Biology, 2003b (in press).

ACKNOWLEDGMENTS

This work was supported by Laboratory Directed Research and Development funds at Berkeley Lab provided by the Director and by the Atmospheric Radiation Measurement Program, Office of Science, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

